

STOBOCLO® Patient Guide:

What is Stoboclo®?

Stoboclo® (denosumab-bmwo) is a prescription medicine used to:


- Treat osteoporosis (thinning and weakening of bone) in women after menopause (“change of life”) who:
 - Are at high risk for fracture (broken bone)
 - Cannot use another osteoporosis medicine or other osteoporosis medicines did not work well
- Treat to increase bone mass in men with osteoporosis who are at high risk for fracture
- Treat bone loss in men who are at high risk for fracture receiving certain treatments for prostate cancer that has not spread to other parts of the body
- Treat bone loss in women who are at high risk for fracture receiving certain treatments for breast cancer that has not spread to other parts of the body
- Treat osteoporosis in men and women who will be taking corticosteroid medicines (such as prednisone) for at least 6 months and are at high risk for fracture

What is the serious risk of Stoboclo®?

Patients with advanced chronic kidney disease, including dialysis-dependent patients, are at risk of severe hypocalcemia (low calcium levels in your blood) following Stoboclo® administration.

Most people with low blood calcium levels do not have symptoms, but some people may have symptoms which include:

- Spasms, twitches, or cramps in your muscles
- Numbness or tingling in your fingers, toes, or around your mouth



Call your doctor right away if you think you may be having any of these symptoms.

What to expect before and during Stoboclo® treatment:

- Talk to your doctor before starting Stoboclo®.
- If you have low blood calcium before you start receiving Stoboclo®, it may get worse during treatment.
- Your low blood calcium must be treated before you receive Stoboclo®.
- Your doctor may prescribe calcium and vitamin D to help prevent low calcium levels in your blood while you take Stoboclo®. Take calcium and vitamin D as your doctor tells you to.
- If you have advanced chronic kidney disease (may or may not be on kidney dialysis), Stoboclo® may increase your risk for severe low calcium levels in your blood, leading to hospitalization, life-threatening events, and death.
- A mineral and bone disorder associated with kidney disease called chronic kidney disease-mineral bone disorder (CKD-MBD) may increase your risk for severe low calcium levels in blood. Before you start Stoboclo® and during treatment, your doctor may need to do certain blood tests to check for CKD-MBD.

If you have advanced chronic kidney disease, your doctor may coordinate your Stoboclo® treatment with a healthcare provider with expertise in CKD-MBD.